

dozes of 25, 50, and 250 mg/kg per day (0.1, 0.3, and 2 times the recommended human dose, respectively, based on body surface area) were used. In the 2-year carcinogenicity study, there was a statistically significant increase in the incidence of benign adrenal pheochromocytoma in male rats treated with the 250 mg/kg per day (about 2 times the maximum recommended human dose, based on body surface area) dose, as compared to vehicle treated rats. The mouse carcinogenicity study showed no evidence of a carcinogenic effect, but the study was not ideal due to its duration of 78 instead of 104 weeks.

Saproterin dihydrochloride was genotoxic in the *in vivo* Ames test at concentrations of 625 mg/kg (7490) and 5,000 mg/kg (74100) per plate, without metabolic activation. However, no genotoxicity was observed in the *in vitro* Ames test with metabolic activation. Saproterin dihydrochloride was genotoxic in the *in vivo* chromosomal aberration assay in Chinese hamster lung cells at concentrations of 0.25 and 0.5 mM. Saproterin dihydrochloride was not mutagenic in the *in vivo* micronucleus assay in mice at doses up to 2,000 mg/kg per day (about 8 times the maximum recommended human dose of 20 mg/kg per day, based on body surface area). Saproterin dihydrochloride, at oral doses up to 400 mg/kg per day (about 3 times the maximum recommended human dose, based on body surface area) was found to have no effect on fertility and reproductive function of male and female rats.

14 CLINICAL STUDIES

The efficacy of saproterin dihydrochloride was evaluated in five clinical studies in patients with PKU.

Study 1 was a multicenter, open-label, uncontrolled clinical trial of 489 patients with PKU, ages 8 to 48 years (mean 22 years), who had baseline blood Phe levels 4500 micromol/L and who were not on Phe-restricted diets. All patients received treatment with saproterin dihydrochloride 10 mg/kg per day for 8 days. For the purposes of this study, response to saproterin dihydrochloride treatment was defined as a $\geq 30\%$ decrease in blood Phe from baseline. At Day 8, 96 patients (20%) were identified as responders.

Study 2 was a multicenter, double-blind, placebo-controlled study of 88 patients with PKU who responded to saproterin dihydrochloride in Study 1. After a washout period from Study 1, patients were randomized equally to either saproterin dihydrochloride 10 mg/kg per day (N=41) or placebo (N=47) for 6 weeks. Efficacy was assessed by the mean change in blood Phe level from baseline to Week 6 in the saproterin dihydrochloride-treated group as compared to the mean change in the placebo group.

The results showed that at baseline, the mean (\pm SD) blood Phe level was 843 (± 300) micromol/L in the saproterin dihydrochloride-treated group and 888 (± 323) micromol/L in the placebo group. At Week 6, the saproterin dihydrochloride-treated group had a mean (\pm SD) blood Phe level of 607 (± 377) micromol/L, and the placebo group had a mean blood Phe level of 891 (± 348) micromol/L. At Week 6, the saproterin dihydrochloride- and placebo-treated groups had mean changes in blood Phe level of -239 and 6 micromol/L, respectively (mean percent changes of -29% (± 32) and 3% (± 33), respectively). The difference between the groups was statistically significant ($p < 0.0001$) (Table 6).


Table 6: Blood Phe Results in Study 2

	Saproterin (N=41)	Placebo (N=47)
Baseline Blood Phe Level* (micromol/L)		
Mean (\pm SD)	843 (± 300)	888 (± 323)
Percentiles (25 th , 75 th)	620, 990	618, 1141
Week 6 Blood Phe Level (micromol/L)		
Mean (\pm SD)	607 (± 377)	891 (± 348)
Percentiles (25 th , 75 th)	307, 812	619, 1143
Mean Change in Blood Phe From Baseline to Week 6 (micromol/L)		
Adjusted Mean (\pm SE)	-239 (± 38)	6 (± 36)
Percentiles (25 th , 75 th)	-397, -92	-96, 93
Mean Percent Change in Blood Phe From Baseline to Week 6		
Mean (\pm SD)	-29 (± 32)	3 (± 33)
Percentiles (25 th , 75 th)	-61, -11	-13, 12

*The mean baseline levels shown in this table represent the mean of 3 pretreatment levels (Wk -2, Wk -1, and Wk 0). Treatment with saproterin dihydrochloride for 6 weeks of forced dose-titration with 3 different doses of saproterin dihydrochloride. Treatments consisted of 2 consecutive 2-week courses of saproterin dihydrochloride at doses of 5, then 20, and then 10 mg/kg per day. Blood Phe level was monitored after 2 weeks of treatment at each dose level. At baseline, mean (\pm SD) blood Phe was 844 (± 398) micromol/L. At the end of treatment with 5, 10, and 20 mg/kg per day, mean (\pm SD) blood Phe levels were 714 (± 380) micromol/L, 640 (± 382) micromol/L, and 581 (± 399) micromol/L, respectively (Table 7).

Change in blood Phe was noted in the saproterin dihydrochloride-treated group at Week 1 and was sustained through Week 6 (Figure 2).

Figure 2: Mean Blood Phenylalanine (Phe) Level Over Time*



*Error bars indicate 95% confidence interval.

Study 3 was a multicenter, open-label, extension study in which 80 patients who responded to saproterin dihydrochloride treatment in Study 1 and completed Study 2 underwent 6 weeks of forced dose-titration with 3 different doses of saproterin dihydrochloride. Treatments consisted of 2 consecutive 2-week courses of saproterin dihydrochloride at doses of 5, then 20, and then 10 mg/kg per day. Blood Phe level was monitored after 2 weeks of treatment at each dose level. At baseline, mean (\pm SD) blood Phe was 844 (± 398) micromol/L. At the end of treatment with 5, 10, and 20 mg/kg per day, mean (\pm SD) blood Phe levels were 714 (± 380) micromol/L, 640 (± 382) micromol/L, and 581 (± 399) micromol/L, respectively (Table 7).

Table 7: Blood Phe Results From Forced Dose-Titration in Study 3

Saproterin Dihydrochloride Dose Level (mg/kg per day)	No. of Patients	Mean (\pm SD) Blood Phe Level (micromol/L)	Mean Changes (\pm SD) in Blood Phe Level From Week 0 (micromol/L)
Baseline (No Treatment)	80	844 (± 398)	—
5	80	744 (± 284)	-100 (± 295)
10	80	640 (± 382)	-204 (± 303)
20	80	581 (± 399)	-263 (± 318)

Study 4 was a multicenter study of 90 pediatric patients with PKU, ages 4 to 12 years, who were on Phe-restricted diets and who had blood Phe levels ≥ 4500 micromol/L at screening. All patients were treated with open-label saproterin dihydrochloride 20 mg/kg per day for 8 days. Response to saproterin dihydrochloride was defined as a $\geq 30\%$ decrease in blood Phe from baseline at Day 8. At Day 8, 50 patients (56%) had a $\geq 30\%$ decrease in blood Phe.

Study 5 was an open label, single arm, multicenter trial in 93 pediatric patients with PKU, aged 1 month to 6 years, who had Phe levels greater than or equal to 350 micromol/L at screening. All patients were treated with saproterin dihydrochloride at 20 mg/kg per day and maintained on a Phe-restricted diet. At Week 4, 57 patients (61%) were identified as responders (defined as $\geq 30\%$ decrease in blood Phe from baseline) (see Figure 1 section 8.4).

16 HOW SUPPLIED/STORAGE AND HANDLING

Saproterin dihydrochloride powder for oral solution is off-white to pale yellow powder packaged in unit dose packets as follows:

- 100 mg saproterin dihydrochloride powder for oral solution per packet:
- Single unit dose packet NDC 43598-477-11
- Carton of 30 unit dose packets NDC 43598-477-30

Store saproterin dihydrochloride powder for oral solution at 20°C to 25°C (68°F to 77°F) [see USP Controlled Room Temperature]. Protect from moisture.

17 PATIENT COUNSELING INFORMATION

Advise the patient or caregiver to read the FDA-approved patient labeling (Patient Information and Instructions for Use).

Hypersensitivity Reactions, Including Anaphylaxis
Advise patients and caregivers to discontinue saproterin dihydrochloride and contact the patient's healthcare provider immediately if they experience symptoms of anaphylaxis, including (but not limited to) wheezing, dyspnea, coughing, hypotension, flushing, nausea, and rash. Continue nutritional management including dietary protein and Phe restriction [see *Warnings and Precautions* (5.1)].

Lower Gastrointestinal Mucosal Inflammation
Advise patients and caregivers to contact their healthcare provider if the patient experiences signs and symptoms suggestive of upper GI mucosal inflammation, including nausea, vomiting, dysphagia, dyspepsia, loss of appetite, oropharyngeal, esophageal, or upper abdominal pain [see *Warnings and Precautions* (5.3)].

Hypophenylalaninemia, [see *Warnings and Precautions* (5.3)]
Advise patients and caregivers that saproterin dihydrochloride may cause hypophenylalaninemia (low blood Phe levels), especially in pediatric patients younger than 7 years of age.

Monitoring of Blood Phe Levels [see *Warnings and Precautions* (5.4)]
Advise patients and caregivers that frequent blood Phe monitoring is important to ensure blood Phe levels are in the desirable range and that they should maintain dietary protein and Phe restriction while on saproterin dihydrochloride. Prolonged hypophenylalaninemia (high blood Phe levels) in patients with PKU can result in severe neurologic damage, including intellectual disability, developmental delay, microcephaly, delayed speech, seizures, and behavioral abnormalities.

Lack of Biochemical Response to Saproterin Dihydrochloride
Some patients do not show a biochemical response (blood Phe reduction) when treated with saproterin dihydrochloride. Advise patients and caregivers to discontinue treatment with saproterin dihydrochloride if the patient does not show an adequate biochemical response in blood Phe after one month of treatment with saproterin dihydrochloride 20 mg/kg per day [see *Dosage and Administration* (2.1), *Warnings and Precautions* (5.4)].

Interaction with Levodopa
Advise patients and caregivers that patients with underlying neurological disorders taking saproterin dihydrochloride in combination with levodopa may experience seizures, exacerbation of seizures, over-stimulation or irritability. Inform patients and caregivers to contact their healthcare provider if the patient has a change in neurologic status during treatment with saproterin dihydrochloride [see *Warnings and Precautions* (5.3)].

Hypersensitivity
Advise patients and caregivers that saproterin dihydrochloride may cause hypersensitivity and to contact their healthcare provider if the patient experiences hypersensitivity, restlessness, fidgeting, or excessive talking [see *Warnings and Precautions* (5.6)].

Dosing and Monitoring [see *Dosage and Administration* (2.1)]
Advise patients and caregivers of the following:

- Saproterin dihydrochloride powder for oral solution should be used in conjunction with a PKU-specific diet, including dietary protein and Phe restriction.
- Dietary protein and Phe intake should not be modified during when assessing biochemical response.
- The patient must be evaluated for changes in blood Phe after being treated with saproterin dihydrochloride at the recommended dose(s) for age to determine if they have a biochemical response and that blood Phe levels and dietary Phe intake should be assessed frequently during the first month of treatment with saproterin dihydrochloride treatment.
- Monitoring of blood Phe levels is important during saproterin dihydrochloride treatment.

Preparation and Administration [see *Dosage and Administration* (2.2)]
Advise patients and caregivers:

- Saproterin dihydrochloride powder for oral solution should be dissolved in water or apple juice or stirred in a small amount of soft food such as apple sauce or pudding.
- Take saproterin dihydrochloride powder for oral solution with a meal, preferably at the same time each day.

Rx only

Distributor:
Dr. Reddy's Laboratories Inc.,
Princeton, NJ 08540
Made in India

Dr.Reddy's

Issued: 1220

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

PATIENT INFORMATION

Saproterin (SAP-ro-TER-in) Dihydrochloride Powder for Oral Solution

What is saproterin dihydrochloride?
Saproterin dihydrochloride powder for oral solution is a prescription medicine used to lower blood levels of phenylalanine (Phe), in adults and children one month of age and older with a certain type of Phenylketonuria (PKU). Saproterin dihydrochloride powder for oral solution is used along with a Phe-restricted diet.

What should I tell my doctor before taking saproterin dihydrochloride powder for oral solution?
Before you take saproterin dihydrochloride powder for oral solution, tell your doctor about all your medical conditions, including if you:

- are allergic to saproterin dihydrochloride or any of the ingredients in saproterin dihydrochloride powder for oral solution. See the list of ingredients in saproterin dihydrochloride powder for oral solution at the end of this leaflet.
- have poor nutrition or have loss of appetite.
- are pregnant or plan to become pregnant.
- are breastfeeding or plan to breastfeed. It is not known if saproterin dihydrochloride passes into your breast milk. Talk to your doctor about the best way to feed your baby if you take saproterin dihydrochloride powder for oral solution.

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, herbal, and dietary supplements. Saproterin dihydrochloride powder for oral solution and other medicines may interact with each other.

Especially tell your doctor if you take:

- a medicine that contains levodopa
- an antileite medicine
- sildenafil (Revatio, Viagra), tadalafil (Addicra, Cialis), vardenafil (Stavyn, Levitra)

Tell your doctor if you are not sure if your medicine is one that is listed above.

Know the medicines you take. Keep a list of them to show your doctor and pharmacist when you get a new medicine.

How should I take saproterin dihydrochloride powder for oral solution?
Take saproterin dihydrochloride powder for oral solution exactly as your doctor tells you. Your doctor should tell you how much saproterin dihydrochloride powder for oral solution to take and when to take it.

Your doctor may change your dose of saproterin dihydrochloride powder for oral solution depending on how you respond to treatment.

Take saproterin dihydrochloride powder for oral solution 1 time each day with a meal. It is best to take saproterin dihydrochloride powder for oral solution at the same time each day.

Saproterin dihydrochloride comes as a powder for oral solution.

Be sure that you know what dose of saproterin dihydrochloride powder your doctor prescribed and whether you should use saproterin dihydrochloride powder for oral solution 100 mg packets, saproterin dihydrochloride powder for oral solution 500 mg packets, or both types of packets to prepare your dose.

Open saproterin dihydrochloride powder packets only when you are ready to use them.

Saproterin dihydrochloride powder for oral solution should be dissolved in water or apple juice. You may also mix the powder for oral solution in a small amount of soft food, such as apple sauce or pudding before taking.

See the detailed "Instructions for Use" that comes with saproterin dihydrochloride powder for oral solution for information about the correct way to dissolve and take a dose of saproterin dihydrochloride powder for oral solution.

It is not possible to know if saproterin dihydrochloride powder for oral solution will work for you until you start taking saproterin dihydrochloride powder for oral solution. Your doctor will check your blood Phe levels when you start taking saproterin dihydrochloride powder for oral solution to see if the medicine is working.

During treatment with saproterin dihydrochloride powder for oral solution:

- Any change you make to your diet may affect your blood Phe level. Follow your doctor's instructions carefully and do not make any changes to your dietary Phe intake without first talking with your doctor. Even if you take saproterin dihydrochloride powder for oral solution, if your Phe blood levels are not well controlled, you can develop severe neurologic problems.
- Your doctor should continue to monitor your blood Phe levels during your treatment with saproterin dihydrochloride, to make sure that your blood Phe levels are not too high or too low.

If you have a fever, or if you are sick, your blood Phe level may go up. Tell your doctor as soon as possible so they can change your dose of saproterin dihydrochloride to help keep your blood Phe level in the desired range.

If you forget to take your dose of saproterin dihydrochloride powder for oral solution, take it as soon as you remember that day. Do not take 2 doses in a day.

If you take too much saproterin dihydrochloride powder for oral solution, call your doctor for advice.

What are the possible side effects of saproterin dihydrochloride powder for oral solution?
Saproterin dihydrochloride powder for oral solution can cause serious side effects, including:

Severe allergic reactions. Stop taking saproterin dihydrochloride powder for oral solution and get medical help right away if you develop any of these symptoms of a severe allergic reaction:

- hives or trouble breathing
- flushing
- coughing
- nausea
- feeling lightheaded or you faint
- rash

Inflammation of the lining of the stomach (gastritis) or esophagus (esophagitis). Gastritis or esophagitis can happen with saproterin dihydrochloride and may be severe. Call your doctor right away if you have any of these signs or symptoms:

- severe upper stomach-area (abdominal) discomfort or pain, nausea and vomiting
- blood in your vomit or stool
- black, tarry stools
- difficulty swallowing
- loss of appetite
- pain in the throat

Phe levels that are too low. Some children under the age of 7 years who take high doses of saproterin dihydrochloride each day may experience low Phe levels.

Toxic shock or constant activity (hyperactivity) can happen with saproterin dihydrochloride. Tell your doctor if you have any signs of hyperactivity, including:

- fidgeting or moving around too much
- talking too much

The most common side effects of saproterin dihydrochloride powder for oral solution are:

- headache
- runny nose and nasal congestion
- sore throat
- diarrhea
- vomiting
- cough

Tell your doctor if you have any side effect that bothers you or that does not go away.

These are not all the possible side effects of saproterin dihydrochloride powder for oral solution. For more information, ask your doctor or pharmacist. Call your doctor about any prescription or over-the-counter medicines you are taking. You may report side effects to FDA at 1-800-FDA-1088.

How should I store saproterin dihydrochloride powder for oral solution?

- Store saproterin dihydrochloride powder for oral solution at room temperature between 20°C to 25°C (68°F to 77°F).
- Protect from moisture.

Keep saproterin dihydrochloride powder for oral solution and all medicines out of the reach of children.

General information about the safe and effective use of saproterin dihydrochloride powder for oral solution.
Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use saproterin dihydrochloride powder for oral solution for a condition for which it was not prescribed. Do not give saproterin dihydrochloride powder for oral solution to other people, even if they have the same symptoms you have. It may harm them. You can ask your pharmacist or doctor for information about saproterin dihydrochloride powder for oral solution that is written for health professionals.

What are the ingredients in saproterin dihydrochloride powder for oral solution?
Active ingredient: saproterin dihydrochloride.
Inactive ingredients: ascorbic acid, mannitol, sucralose and tripotassium citrate monohydrate.

For more information, call 1-888-375-3784.

This Patient Information has been approved by the U.S. Food and Drug Administration.
To reorder additional Patient Information Sheets contact Dr. Reddy's Customer Service at 1-866-733-3952.

Rx only

Distributor:
Dr. Reddy's Laboratories Inc.,
Princeton, NJ 08540
Made in India

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's

Dr.Reddy's